


**KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING**



PERMETHRIN 25:75 INSECTICIDAL DUSTING POWDER INDUSTRIAL STRENGTH

Active Constituent: 10g/kg PERMETHRIN 25:75

GROUP 3A INSECTICIDE

**For the control of Cockroaches, Ants, Fleas, Termites
European Wasps and other insects in certain situations as per
the Directions for Use**

D.O.M.

CONTENTS: **

Batch no:

NRA Approval no: **55091/0202**

A & C Rural Pty Ltd
A.B.N. 52 010 865 317
Lot 96 Potassium Street
Narangba Qld 4503
Ph: (07) 3203 3944

** This label is to be used for 1kg, 2kg, 2.5kg, 5kg, 10kg, 15kg, 20kg pack sizes.

GENERAL INSTRUCTIONS

Insecticide Resistance Warning

For insecticide resistance management, Country Permethrin 25:75 Insecticidal Dusting Powder Insecticide is a Group 3A insecticide. Some naturally occurring insect biotypes resistant to Country Permethrin 25:75 Insecticidal Dusting Powder and other Group 3A insecticides may exist through normal genetic variability in any insect population. The resistant individuals can eventually dominate the insect population if Country Insecticidal Dusting Powder or other Group 3A insecticides are used repeatedly. The effectiveness of Country Insecticidal Dusting Powder on resistant individuals could be significantly reduced. Since the occurrence of resistant individuals is difficult to detect prior to use, A & C Rural Pty Ltd accepts no liability for any losses that may result from the failure of Country Insecticidal Dusting Powder to control resistant insects. Country Insecticidal Dusting Powder may be subject to specific resistance management strategies. For further information, contact your local supplier or A & C Rural Pty Ltd representative.

APPLICATION

Country Insecticidal Dusting Powder is ready to use; no further diluent should be added. Leave powder undisturbed for as long as possible after application.

PRECAUTIONS

Avoid contact with food, food utensils, or places where food is prepared or stored. Before use, remove or cover all exposed foodstuffs. Cover all dishes and utensils, and places where food is prepared or stored. Do not apply dust directly on humans. Remove or cover fish bowls during treatment.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

Dangerous to fish and other aquatic organisms. DO NOT contaminate streams, rivers or waterways with the chemical or used containers. Dangerous to bees and reptiles.

STORAGE AND DISPOSAL

For plastic tubs: Keep out of reach of children. Store in the closed, original container in a cool, well-ventilated area. Do not store for prolonged periods in direct sunlight. Triple or preferably pressure rinse containers before disposal. Dispose of rinsings in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Break, crush or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

SAFETY DIRECTIONS

Will irritate the eyes. Avoid contact with eyes. Do not inhale dust. When opening the container and using the product, wear rubber gloves and a disposable dust mask. Wash hands after use. After each days use, wash gloves and contaminated clothing.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre (phone 13 11 26).

MATERIAL SAFETY DATA SHEET

Additional information is listed in the Material Safety Data Sheet which is available from A & C Rural Pty Ltd upon request.

CONDITIONS OF SALE

The use of Country Permethrin Insecticidal Dusting Powder Insecticide being beyond the control of the manufacturer no warranty expressed or implied is given by A & C Rural Pty Ltd regarding its suitability, fitness or efficiency for any purpose for which it is used by the buyer, whether in accordance with the directions or not and A & C Rural Pty Ltd accepts no responsibility for any consequence whatsoever resulting from the use of this product.

DIRECTIONS FOR USE:

SITUATION	PEST	RATE	CRITICAL COMMENTS
Domestic, institutional and industrial premises - food preparation and storage areas and living and sleeping areas	Cockroaches, silverfish	10g/m ² applied with a powder duster	Blow powder into all cracks and crevices. Pay particular attention to warm, dark corners around cookers, fireplaces, radiators, stores, pipes, rubbish chutes and places where liquid sprays may not be used.
	Ants		Sprinkle the powder along runs, on and around nests. If ants are invading domestic premises, apply powder to the routes by which they enter. Repeat as necessary.
	Fleas		Dust powder over floor, brushing into carpet pile and blowing into areas where flea larvae develop.
	Bedbugs		Apply powder to bedsteads, bedsprings, mattresses, floor coverings, upholstered furniture, cracks in walls, behind torn wallpaper, joints in woodwork and other cracks and crevices that may provide harbourages.
Roof voids, wall and floor voids, crawl spaces	Spiders, cockroaches, silverfish, ants and fleas	125g to 250g per nest	Apply dust into all void areas. Ensure all surfaces are treated. Repeat as necessary if re-infestation occurs.
	Bird mites (<i>Ornithonyssus spp</i>)		Cover roof void area, paying particular attention to areas where there is evidence of bird nests. Remove nests, and bird-proof roof voids to prevent recurrence.
Skin and hide preparation	Hide Beetles (<i>Dermestes ater</i>)		a) Place skins on drying racks and apply dust using a manual or motorised duster. b) Dust exposed edges of dried, baled skins.
Mushroom beds	Mushroom sciarid flies		Dust mushroom beds no more frequently than twice a week to control adult and larval flies. Discontinue dusting when mushrooms reach the size of a 10-cent piece.
Large wasp nests	European wasps <u>Not Qld, SA</u>	125g to 250g per nest	Using hand duster apply powder through entrance holes (125-250g per nest depending on size of nest and number of entrance holes). Treat at night whenever possible. Operators should wear clothing protective against the wasp including long-sleeved overalls buttoned at wrist and throat, gloves, hat and bee-veil.
Seed potatoes	Potato Moth (<i>Phthorimaea operculella</i>)	600g per tonne or 30g per 50kg bag of seed potatoes	Dust thoroughly immediately after harvest while filling storage containers. Treated seed potatoes are to be used only for sowing. Do not use for food or feed purposes. Treated seed potatoes should be stored apart from other potatoes and bags or containers should be clearly marked to indicate that the contents have been treated.
Trees and stumps in urban areas	Subterranean termites (<i>Coptotermes spp</i>)	250g per tree	Upon locating nest in tree or stump, drill 3 holes with a maximum diameter of 12mm. Apply dust directly to the nest using a carbon dioxide gas cylinder and a trigger gun. After treatment, holes should be coated carefully with a fungicide, e.g. copper naphthenate, and sealed with an appropriate caulking compound.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIOD:

Mushrooms: DO NOT HARVEST FOR 3 DAYS AFTER APPLICATION